

## ARK™ Voriconazole II Assay


The ARK™ Voriconazole II Assay is intended for the quantitative determination of voriconazole in human serum on automated clinical chemistry analyzers. The measurements obtained are used in monitoring levels of voriconazole to help ensure appropriate therapy.




### KEY POINTS

- Homogeneous Enzyme immunoassay
- Applicable onboard automated clinical chemistry analyzers
- Convenient, liquid-stable, ready-to-use
- Excellent calibration range
- Tested drugs and endogenous substances do not interfere
- The crossreactivity of N-oxide voriconazole metabolite was not clinically significant ( $\leq 3.0\%$ )

*Next Generation Assays*

**CALIBRATION RANGE**

ARK™ Voriconazole II Assay Calibration Range: 0.0 to 16.0 µg/mL.  
LOQ: 0.5 µg/mL

**METHOD COMPARISON**

ARK™ Voriconazole II Assay: Measurement Range: 0.5 to 14.0 µg/mL.

**PRECISION**

Sample	N	WITHIN-RUN			BETWEEN DAY			TOTAL	
		Mean (µg/mL)	SD	CV (%)	SD	CV (%)	SD	CV (%)	
<b>ARK Voriconazole II Control</b>									
LOW	160	1.03	0.047	4.6	0.022	2.1	0.051	4.9	
MID	160	4.91	0.194	3.9	0.101	2.1	0.209	4.3	
HIGH	160	9.39	0.394	4.2	0.207	2.2	0.426	4.5	
<b>Human Serum</b>									
LOW	160	1.02	0.043	4.2	0.024	2.4	0.047	4.6	
MID	160	5.03	0.182	3.6	0.111	2.2	0.217	4.3	
HIGH	160	9.80	0.334	3.4	0.221	2.3	0.407	4.2	

Tri-level controls and sera containing voriconazole were assayed in quadruplicate twice a day for 20 days. CLSI/NCCLS Protocol EP5-A3.

**ACCURACY**

Theoretical Concentration (µg/mL)	Mean Recovered Concentration (µg/mL)	Percent Recovery
0.5	0.45	90.0
1.2	1.19	99.2
3.0	3.05	101.7
6.0	5.86	97.7
9.0	8.74	97.1
12.0	11.44	95.3
15.0	15.75	105.0

Accuracy (analytical recovery) was determined by adding concentrated voriconazole drug into human serum negative to produce across the assay range.

**INTERFERENCE**

Tested endogenous substances and co-administered drugs do not interfere with ARK™ Voriconazole II Assay. The crossreactivity of major metabolite N-oxide voriconazole was not clinically significant ( $\leq 3.0\%$  crossreactivity).

**SAFETY AND STABILITY****Reagent on-board stability**

At least 60 days

**Shelf Life of Reagents, Calibrators, and Controls**

18 months from date of manufacturing

**Safety**

Nonhazardous preservatives  
Contains sodium azide  $\leq 0.09\%$

Results shown are typical and may vary among laboratory analyzers.

Available Upon request: UK NEQAS Proficiency Testing for Antifungal Drugs.

**ORDERING INFORMATION**

ARK™ Voriconazole II Assay 5030-0001-01

ARK™ Voriconazole II Calibrator 5030-0002-01

ARK™ Voriconazole II Control 5030-0003-01

**ARK Diagnostics, Inc.**

48089 Fremont Boulevard

Fremont, CA 94538

Tel: 510-270-6270 Fax: 510-270-6298

**For Customer Support:**

Call toll free: 877-869-2320

customersupport@ark-tdm.com

[www.ark-tdm.com](http://www.ark-tdm.com)



**ARK** Diagnostics, Inc.

© 2017 ARK Diagnostics, Inc. All rights reserved  
All trademarks are property of ARK Diagnostics, Inc.

MKT15-016 Rev 02 May 2017